

## SEMI-INVARIANTS OF QUIVERS AND SATURATION FOR LITTLEWOOD-RICHARDSON COEFFICIENTS

HARM DERKSEN AND JERZY WEYMAN

### 1. INTRODUCTION

Let  $Q$  be a quiver without oriented cycles. Let  $\alpha$  be a dimension vector for  $Q$ . We denote by  $\text{SI}(Q, \alpha)$  the ring of semi-invariants of the set of  $\alpha$ -dimensional representations of  $Q$  over a fixed algebraically closed field  $K$ .

In this paper we prove some results about the set

$$\Sigma(Q, \alpha) = \{ \sigma \mid \text{SI}(Q, \alpha)_\sigma \neq 0 \}.$$

$\Sigma(Q, \alpha)$  is defined in the space of all weights by one homogeneous linear equation and by a finite set of homogeneous linear inequalities. In particular the set  $\Sigma(Q, \alpha)$  is saturated, i.e., if  $n\sigma \in \Sigma(Q, \alpha)$ , then also  $\sigma \in \Sigma(Q, \alpha)$ .

These results, when applied to a special quiver  $Q = T_{n,n,n}$  and to a special dimension vector, show that the  $\text{GL}_n$ -module  $V_\lambda$  appears in  $V_\mu \otimes V_\nu$  if and only if the partitions  $\lambda$ ,  $\mu$  and  $\nu$  satisfy an explicit set of inequalities. This gives new proofs of the results of Klyachko ([7, 3]) and Knutson and Tao ([8]).

The proof is based on another general result about semi-invariants of quivers (Theorem 1). In the paper [10], Schofield defined a semi-invariant  $c_W$  for each indecomposable representation  $W$  of  $Q$ . We show that the semi-invariants of this type span each weight space in  $\text{SI}(Q, \alpha)$ . This seems to be a fundamental fact, connecting semi-invariants and modules in a direct way. Given this fact, the results on sets of weights follow at once from the results in another paper of Schofield [11].

### 2. THE RESULTS

A quiver  $Q$  is a pair  $Q = (Q_0, Q_1)$  consisting of the set of vertices  $Q_0$  and the set of arrows  $Q_1$ . Each arrow  $a$  has its head  $ha$  and tail  $ta$ , both in  $Q_0$ :

$$ta \xrightarrow{a} ha.$$

We fix an algebraically closed field  $K$ . A representation (or a module)  $V$  of  $Q$  is a family of finite dimensional vector spaces  $\{V(x) \mid x \in Q_0\}$  and of linear maps

---

Received by the editors July 20, 1999.

2000 *Mathematics Subject Classification*. Primary 13A50; Secondary 14L24, 14L30, 16G20, 20G05.

*Key words and phrases*. Quiver representations, semi-invariants, Littlewood-Richardson coefficients, Klyachko cone, saturation.

The second author was supported by NSF, grant DMS 9700884 and KBN No. PO3A 012 14.

$V(a) : V(ta) \rightarrow V(ha)$ . The dimension vector of a representation  $V$  is the function  $\underline{d}(V) : Q_0 \rightarrow \mathbb{Z}_{\geq 0}$  defined by  $\underline{d}(V)(x) := \dim V(x)$ . The dimension vectors lie in the space  $\Gamma$  of integer-valued functions on  $Q_0$ . A morphism  $\phi : V \rightarrow V'$  of two representations is a collection of linear maps  $\phi(x) : V(x) \rightarrow V'(x)$ ,  $x \in Q_0$ , such that for each  $a \in Q_1$  we have  $\phi(ha)V(a) = V'(a)\phi(ta)$ . We denote the linear space of morphisms from  $V$  to  $V'$  by  $\text{Hom}_Q(V, V')$ .

A path  $p$  in  $Q$  is a sequence of arrows  $p = a_1, \dots, a_n$  such that  $ha_i = ta_{i+1}$  ( $1 \leq i \leq n - 1$ ). We define  $tp = ta_1, hp = ha_n$ . We also have the trivial path  $e(x)$  from  $x$  to  $x$ . If  $V$  is a representation and  $p = a_1, \dots, a_n$ , then we define  $V(p) := V(a_n)V(a_{n-1}) \cdots V(a_1)$ . We assume throughout the paper that  $Q$  has no oriented cycles, i.e., there are no paths  $p = a_1, \dots, a_n$  such that  $ta_1 = ha_n$ .

For representations  $V$  and  $W$  of  $Q$  there is a canonical exact sequence ([9])

$$(1) \quad 0 \rightarrow \text{Hom}_Q(V, W) \xrightarrow{i} \bigoplus_{x \in Q_0} \text{Hom}(V(x), W(x)) \xrightarrow{d_W^V} \bigoplus_{a \in Q_1} \text{Hom}(V(ta), W(ha)) \xrightarrow{p} \text{Ext}_Q(V, W) \rightarrow 0.$$

The map  $i$  is the obvious inclusion, the map  $d_W^V$  is given by

$$\{f(x)\}_{x \in Q_0} \mapsto \{f(ha)V(a) - W(a)f(ta)\}_{a \in Q_1},$$

and the map  $p$  constructs an extension of the representations  $V$  and  $W$  by adding the maps  $V(ta) \rightarrow W(ha)$  to the direct sum representation  $V \oplus W$ .

For  $\alpha, \beta \in \Gamma$  we define the Euler inner product

$$\langle \alpha, \beta \rangle = \sum_{x \in Q_0} \alpha(x)\beta(x) - \sum_{a \in Q_1} \alpha(ta)\beta(ha).$$

It follows from (1) that  $\langle \underline{d}(V), \underline{d}(W) \rangle = \dim_K \text{Hom}_Q(V, W) - \dim_K \text{Ext}_Q(V, W)$ .

For a dimension vector  $\alpha$  we denote by

$$\text{Rep}(Q, \alpha) := \bigoplus_{a \in Q_1} \text{Hom}(K^{\alpha(ta)}, K^{\alpha(ha)})$$

the vector space of  $\alpha$ -dimensional representations of  $Q$ . The group

$$\text{GL}(Q, \alpha) := \prod_{x \in Q_0} \text{GL}(\alpha(x))$$

and its subgroup

$$\text{SL}(Q, \alpha) = \prod_{x \in Q_0} \text{SL}(\alpha(x))$$

act on  $\text{Rep}(Q, \alpha)$  in an obvious way. We are interested in the ring of semi-invariants

$$\text{SI}(Q, \alpha) := K[\text{Rep}(Q, \alpha)]^{\text{SL}(Q, \alpha)}.$$

The ring  $\text{SI}(Q, \alpha)$  has a weight space decomposition

$$\text{SI}(Q, \alpha) = \bigoplus_{\sigma} \text{SI}(Q, \alpha)_{\sigma}$$

where  $\sigma$  runs through the (one-dimensional irreducible) characters of  $\text{GL}(Q, \alpha)$  and

$$\text{SI}(Q, \alpha)_{\sigma} = \{ f \in K[\text{Rep}(Q, \alpha)] \mid g(f) = \sigma(g)f \ \forall g \in \text{GL}(Q, \alpha) \}.$$

Suppose that  $\sigma$  lies in the dual space  $\Gamma^* := \text{Hom}(\Gamma, \mathbb{Z})$ . For each dimension vector  $\alpha$  we can associate to  $\sigma$  a character of  $\text{GL}(Q, \alpha)$  defined as

$$\prod_{x \in Q_0} d_x^{\sigma(e_x)}$$

where  $d_x$  is the determinant function on  $\text{GL}(\alpha(x))$  and  $e_x$  is the dimension vector defined by

$$e_x(y) = \begin{cases} 1 & \text{if } x = y, \\ 0 & \text{otherwise.} \end{cases}$$

In this way we will identify characters with  $\Gamma^*$ . Sometimes, for convenience, we will write  $\sigma(x)$  instead of  $\sigma(e_x)$  (and treat  $\sigma$  as an element of  $\Gamma$ ).

Let us choose the dimension vectors  $\alpha$  and  $\beta$  in such way that  $\langle \alpha, \beta \rangle = 0$ . Then for every  $V \in \text{Rep}(Q, \alpha)$  and  $W \in \text{Rep}(Q, \beta)$  the matrix of  $d_W^V$  will be a square matrix. Following [10] we can therefore define the semi-invariant  $c$  of the action of  $\text{GL}(Q, \alpha) \times \text{GL}(Q, \beta)$  on  $\text{Rep}(Q, \alpha) \times \text{Rep}(Q, \beta)$  by  $c(V, W) := \det d_W^V$ . The value of the determinant depends on the choices of bases, so  $c$  is well-defined up to a scalar. Notice that the semi-invariant  $c$  vanishes at the point  $(V, W)$  if and only if  $\text{Hom}_Q(V, W) \neq 0$  which is equivalent to  $\text{Ext}_Q(V, W) \neq 0$ . For a fixed  $V$  the restriction of  $c$  to  $\{V\} \times \text{Rep}(Q, \beta)$  defines a semi-invariant  $c^V$  in  $\text{SI}(Q, \beta)$ . Schofield proves ([10, Lemma 1.4]) that the weight of  $c^V$  equals  $\langle \alpha, \cdot \rangle \in \Gamma^*$  which is defined as  $\gamma \mapsto \langle \alpha, \gamma \rangle$ . Similarly, for a fixed  $W$  the restriction of  $c$  to  $\text{Rep}(Q, \alpha) \times \{W\}$  defines a semi-invariant  $c_W$  in  $\text{SI}(Q, \alpha)$  of weight  $-\langle \cdot, \beta \rangle$  ([10, Lemma 1.4]). If  $V, V' \in \text{Rep}(Q, \alpha)$  and  $V \cong V'$ , then  $V$  and  $V'$  are in the same  $\text{GL}(Q, \alpha)$ -orbit, and  $c^V$  and  $c^{V'}$  are equal up to a constant scalar. Semi-invariants of the types  $c^V$  and  $c_W$  are well-defined up to a scalar. These semi-invariants have the following properties.

**Lemma 1.** *Suppose that  $V, V', V''$  and  $W, W', W''$  are representations of  $Q$  such that  $\langle \underline{d}(V), \underline{d}(W) \rangle = 0$ , and that there are exact sequences*

$$0 \rightarrow V' \rightarrow V \rightarrow V'' \rightarrow 0, \quad 0 \rightarrow W' \rightarrow W \rightarrow W'' \rightarrow 0.$$

- a) *If  $\langle \underline{d}(V'), \underline{d}(W) \rangle < 0$ , then  $c^V(W) = 0$ ;*
- b) *If  $\langle \underline{d}(V'), \underline{d}(W) \rangle = 0$ , then  $c^V(W) = c^{V'}(W)c^{V''}(W)$ ;*
- c) *If  $\langle \underline{d}(V), \underline{d}(W') \rangle > 0$ , then  $c^V(W) = 0$ ;*
- d) *If  $\langle \underline{d}(V), \underline{d}(W') \rangle = 0$ , then  $c^V(W) = c^V(W')c^V(W'')$ .*

*Proof.* Consider the following commutative diagram with exact columns:

$$\begin{array}{ccc}
 0 & & 0 \\
 \downarrow & & \downarrow \\
 \bigoplus_{x \in Q_0} \text{Hom}(V''(x), W(x)) & \xrightarrow{d_W^{V''}} & \bigoplus_{a \in Q_1} \text{Hom}(V''(ta), W(ha)) \\
 \downarrow & & \downarrow \\
 \bigoplus_{x \in Q_0} \text{Hom}(V(x), W(x)) & \xrightarrow{d_W^V} & \bigoplus_{a \in Q_1} \text{Hom}(V(ta), W(ha)) \\
 \downarrow & & \downarrow \\
 \bigoplus_{x \in Q_0} \text{Hom}(V'(x), W(x)) & \xrightarrow{d_W^{V'}} & \bigoplus_{a \in Q_1} \text{Hom}(V'(ta), W(ha)) \\
 \downarrow & & \downarrow \\
 0 & & 0
 \end{array}$$

If  $\langle \underline{d}(V'), \underline{d}(W) \rangle = 0$ , then  $d_W^{V'}$ ,  $d_W^V$  and  $d_W^{V''}$  are all represented by square matrices. It follows that  $c^V(W) = c^{V'}(W)c^{V''}(W)$ . So b) follows and d) goes similarly. If  $\langle \underline{d}(V'), \underline{d}(W) \rangle < 0$ , then  $d_W^{V'}$  cannot be surjective, hence  $d_W^V$  is not surjective. Now a) follows and c) goes similarly.  $\square$

Our main result is that the semi-invariants of type  $c^V$  (resp.  $c_W$ ) span all the weight spaces in the rings  $\text{SI}(Q, \alpha)$ .

**Theorem 1.** *Let  $Q$  be a quiver without oriented cycles and let  $\beta$  be a dimension vector. The ring of semi-invariants  $\text{SI}(Q, \beta)$  is a  $K$ -linear span of semi-invariants  $c^V$  with  $\langle \underline{d}(V), \beta \rangle = 0$ . The analogous result is true for the semi-invariants  $c_W$ .*

After this paper was submitted we learned about the paper [12] where among other things the authors give another proof of Theorem 1 under the assumption that the characteristic of  $K$  is zero.

We will prove Theorem 1 in Section 4.

*Remark 1.* If  $V = V_1 \oplus V_2$  is decomposable, then by Lemma 1 we have  $c^V = 0$  if  $\langle \underline{d}(V_1), \beta \rangle \neq 0$ , and  $c^V = c^{V_1}c^{V_2}$  if  $\langle \underline{d}(V_1), \beta \rangle = 0$ .

The algebra  $\text{SI}(Q, \beta)$  is generated by all  $c^V$  where  $V$  is *indecomposable*. Generators of  $\text{SI}(Q, \beta)$  therefore can be found in the degrees  $\langle \alpha, \cdot \rangle$  such that a general representation of dimension  $\alpha$  is indecomposable. By [5] this is equivalent to  $\alpha$  being a Schur root.

*Remark 2.* If  $\text{Rep}(Q, \beta)$  has a dense  $\text{GL}(Q, \beta)$ -orbit, then Schofield showed in [10] that the invariants of type  $c^V$  with  $V$  indecomposable generate  $\text{SI}(Q, \beta)$  (which is a polynomial ring in this case).

Theorem 1 has the following remarkable consequence.

**Corollary 1** (Reciprocity Property). *Let  $\alpha, \beta$  be two dimension vectors for the quiver  $Q$ . Assume that  $\langle \alpha, \beta \rangle = 0$ . Then*

$$\dim_K \text{SI}(Q, \beta)_{\langle \alpha, \cdot \rangle} = \dim_K \text{SI}(Q, \alpha)_{-\langle \cdot, \beta \rangle}.$$

*Proof.* Let  $V_1, \dots, V_s$  be the modules of dimension  $\alpha$  such that  $c^{V_1}, \dots, c^{V_s}$  form a basis of  $\text{SI}(Q, \beta)_{\langle \alpha, \cdot \rangle}$ . These are linearly independent polynomials on  $\text{Rep}(Q, \beta)$  so there exist  $s$  representations  $W_1, \dots, W_s$  in  $\text{Rep}(Q, \beta)$  such that  $\det(c^{V_i}(W_j))_{1 \leq i, j \leq s}$

is not zero. But  $c^{V_i}(W_j) = c_{W_j}(V_i)$  and this means that the semi-invariants  $c_{W_1}, \dots, c_{W_s}$  are linearly independent. This proves that

$$\dim_K \text{SI}(Q, \beta)_{\langle \alpha, \cdot \rangle} \leq \dim_K \text{SI}(Q, \alpha)_{-\langle \cdot, \beta \rangle}.$$

The other inequality is proven in exactly the same way.  $\square$

In the remainder of this section we investigate the consequences of Theorem 1. First we recall the main results of [11]. They can be summarized as follows.

We say that for two dimension vectors  $\alpha, \beta$  the space  $\text{Hom}_Q(\alpha, \beta)$  (respectively  $\text{Ext}_Q(\alpha, \beta)$ ) vanishes generically if and only if for general representations  $V, W$  of dimensions  $\alpha, \beta$  respectively we have  $\text{Hom}_Q(V, W) = 0$  (resp.  $\text{Ext}_Q(V, W) = 0$ ). We also write  $\alpha \hookrightarrow \beta$  if a general representation of dimension  $\beta$  has a subrepresentation of dimension  $\alpha$ .

**Theorem 2** (Schofield). *Let  $\alpha$  and  $\beta$  be two dimension vectors for the quiver  $Q$ .*

- a)  $\text{Ext}_Q(\alpha, \beta)$  vanishes generically if and only if  $\alpha \hookrightarrow \alpha + \beta$ ,
- b)  $\text{Ext}_Q(\alpha, \beta)$  does not vanish generically if and only if  $\beta' \hookrightarrow \beta$  and  $\langle \alpha, \beta - \beta' \rangle < 0$  for some dimension vector  $\beta'$ .

Part a) is proven in Section 3 of [11], and part b) is proven in Section 5.

*Remark 3.* Suppose that  $V$  and  $W$  are general modules of dimension  $\alpha$  and  $\beta$  respectively, such that  $\langle \alpha, \beta \rangle = 0$ . The condition in b) is equivalent to  $\exists \beta' \beta' \hookrightarrow \beta$  such that  $\langle \alpha, \beta' \rangle > 0$ . If  $c^V(W) = 0$ , then  $W$  must have a submodule  $W'$  such that  $\langle \alpha, \underline{d}(W') \rangle > 0$ . This means that the converse of Lemma 1.c) is true for general  $V$  and  $W$ .

**Theorem 3.** *Let  $Q$  be a quiver without oriented cycles and let  $\beta$  be a dimension vector. The semigroup  $\Sigma(Q, \beta)$  is the set of all  $\sigma \in \Gamma$  such that  $\sigma(\beta) = 0$  and  $\sigma(\beta') \leq 0$  for all  $\beta'$  such that  $\beta' \hookrightarrow \beta$ . Thus this condition is provided by one linear homogeneous equality and finitely many linear homogeneous inequalities. In particular the set  $\Sigma(Q, \beta)$  is saturated in the lattice  $\Gamma$ .*

*Proof.* Suppose that  $\sigma \in \Gamma^*$ . We can write  $\sigma = \langle \alpha, \cdot \rangle$  with  $\alpha \in \Gamma$ .

We will first assume that  $\alpha$  is a dimension vector, i.e.,  $\alpha(x) \geq 0$  for all  $x \in Q_0$ . It follows from Theorem 1 that  $\text{SI}(Q, \beta)_{\langle \alpha, \cdot \rangle}$  is non-zero if and only if there exists a representation  $V$  of dimension  $\alpha$  such that  $c^V$  is not zero, which is equivalent to  $\sigma(\beta) = \langle \alpha, \beta \rangle = 0$  and  $\text{Ext}_Q(\alpha, \beta)$  vanishing generically. By part b) of Theorem 2,  $\text{Ext}_Q(\alpha, \beta)$  vanishes generically if and only if for all  $\beta'$  such that  $\beta' \hookrightarrow \beta$  we have  $\langle \alpha, \beta - \beta' \rangle \geq 0$ . This means that for all  $\beta'$  such that  $\beta' \hookrightarrow \beta$  we have  $\sigma(\beta') = \langle \alpha, \beta' \rangle \leq 0$ . We conclude that  $\text{SI}(Q, \beta)_\sigma \neq 0$  if and only if  $\sigma(\beta) = 0$  and  $\sigma(\beta') \leq 0$  for all  $\beta' \hookrightarrow \beta$ .

If  $\alpha$  is not a dimension vector, then  $\text{SI}(Q, \beta)_{n\sigma} = 0$  for all integers  $n > 0$ . Suppose that  $W \in \text{Rep}(Q, \beta)$ . From [6] it follows that either  $\sigma(\underline{d}(W)) \neq 0$  or there exists a submodule  $W'$  of  $W$  such that  $\sigma(\underline{d}(W')) > 0$ . If  $W$  is in general position, then we obtain  $\sigma(\beta) \neq 0$  or  $\sigma(\beta') > 0$  for some  $\beta' \hookrightarrow \beta$  (see also Remark 5).  $\square$

*Remark 4.* Schofield in [11] gives an algorithm allowing one to determine the set of inequalities in Theorem 3 inductively. This algorithm is not very efficient.

*Remark 5.* A module  $W \in \text{Rep}(Q, \beta)$  is called  $\sigma$ -stable if and only if there exist an  $n > 0$  and an  $f \in \text{SI}(Q, \beta)_{n\sigma}$  such that  $f(W) \neq 0$ . King proved in [6] that a module


$W \in \text{Rep}(Q, \beta)$  is  $\sigma$ -stable if and only if  $\sigma(W') \leq 0$  for all submodules  $W'$  of  $W$ . Applied to a general representation  $W$  of dimension  $\beta$  this gives us the equivalence:

$$\exists n > 0 \text{ SI}(Q, \beta)_{n\sigma} \neq 0 \Leftrightarrow \sigma(\beta) = 0 \text{ and } \forall \beta' \beta' \hookrightarrow \beta \text{ we have } \sigma(\beta') \leq 0.$$

This shows that the saturation of  $\Sigma(Q, \beta)$  is given by linear inequalities but it does not show that  $\Sigma(Q, \beta)$  is saturated.

*Remark 6.* In Theorem 3, instead of considering *all*  $\beta'$  with  $\beta' \hookrightarrow \beta$  we only need to consider those  $\beta'$  such that the general representation of dimension  $\beta'$  is indecomposable, which is equivalent to  $\beta'$  being a Schur root. Still, the set of inequalities obtained in this way may not be a minimal set of inequalities as we will see in the next example.

**Example 1.** Let  $Q$  be the quiver



and let  $\beta$  be the dimension vector

$$\begin{array}{c}
 1 \\
 1 \ 2 \ 1 \ . \\
 1
 \end{array}$$

For a general representation  $V$  of  $Q$  with dimension vector  $\beta$ , the dimension vectors of indecomposable submodules are:

$$\begin{array}{cccc}
 0 & 1 & 1 & 1 \\
 1 \ 2 \ 1 & 1 \ 2 \ 0 & 1 \ 2 \ 1 & 0 \ 2 \ 1 \\
 1 & 1 & 0 & 1
 \end{array}$$

$$\begin{array}{cccc}
 1 & 0 & 0 & 0 \\
 0 \ 1 \ 0 & 0 \ 1 \ 1 & 0 \ 1 \ 0 & 1 \ 1 \ 0 \\
 0 & 0 & 1 & 0
 \end{array}$$

$$\begin{array}{c}
 0 \\
 0 \ 1 \ 0 \\
 0
 \end{array}$$

Let  $\sigma$  be the weight given by  $\sigma(\alpha) = \sum_{i=1}^5 a_i \alpha(i)$ , in other words

$$\sigma = \begin{array}{c} a_1 \\ a_4 \ a_5 \ a_2 \ . \\ a_3 \end{array}$$

We investigate when  $\text{SI}(Q, \beta)_\sigma \neq 0$ . First of all we must have  $\sigma(\beta) = 0$ , so  $a_1 + a_2 + a_3 + a_4 + 2a_5 = 0$ . In particular  $a_1 + a_2 + a_3 + a_4$  must be even. The

indecomposable submodules listed above correspond to the inequalities (using  $a_5 = -(a_1 + a_2 + a_3 + a_4)/2$ ):


$$(2) \quad \begin{aligned} & a_1 \geq 0, a_2 \geq 0, a_3 \geq 0, a_4 \geq 0, \\ & a_1 \leq a_2 + a_3 + a_4, a_2 \leq a_1 + a_3 + a_4, a_3 \leq a_1 + a_2 + a_4, a_4 \leq a_1 + a_2 + a_3, \\ & a_1 + a_2 + a_3 + a_4 \geq 0. \end{aligned}$$

The last inequality is redundant.

In the next section we will see how semi-invariants can be interpreted in terms of tensor products of modules of the general linear group. This particular example shows that for a 2-dimensional vector space  $U$ , the tensor product of symmetric powers  $S_{a_1}(U) \otimes S_{a_2}(U) \otimes S_{a_3}(U) \otimes S_{a_4}(U)$  contains a non-trivial  $SL(U)$ -invariant subspace if and only if  $a_1 + a_2 + a_3 + a_4$  is even and the inequalities (2) hold. In this case, the inequalities are obvious from the Clebsch-Gordan formula.

### 3. APPLICATION TO LITTLEWOOD-RICHARDSON COEFFICIENTS

Let us apply Theorem 3 in the following special case. Let us define the quiver  $Q = T_{n,n,n}$  as follows:



Let us choose the dimension vector  $\beta(x_i) = \beta(y_i) = \beta(z_i) = i$  for  $i = 1, \dots, n - 1$ ,  $\beta(u) = n$ . The following proposition is a direct application of Cauchy's formula and is a standard calculation in representation theory.

**Proposition 1.** *The weight space  $SI(T_{n,n,n}, \beta)_\sigma$  is isomorphic to the space of  $SL(U)$ -invariants in the triple tensor product  $S_\lambda(U) \otimes S_\mu(U) \otimes S_\nu(U)$  of Schur functors on  $U$ , where  $U$  is the vector space of dimension  $n$ , and  $\lambda, \mu, \nu$  are partitions whose conjugate partitions are given as follows:*

$$(3) \quad \begin{aligned} \lambda' &= ((n - 1)^{\sigma(x_{n-1})}, (n - 2)^{\sigma(x_{n-2})}, \dots, 1^{\sigma(x_1)}), \\ \mu' &= ((n - 1)^{\sigma(y_{n-1})}, (n - 2)^{\sigma(y_{n-2})}, \dots, 1^{\sigma(y_1)}), \\ \nu' &= ((n - 1)^{\sigma(z_{n-1})}, (n - 2)^{\sigma(z_{n-2})}, \dots, 1^{\sigma(z_1)}). \end{aligned}$$

Here  $\sigma(q)$  is defined as  $\sigma(e_q)$  where the dimension vector  $e_q$  is given by  $e_q(q) = 1$  and  $e_q(p) = 0$  if  $p \neq q$ .

*Proof.* Let us denote by  $a_i$  (resp.  $b_i, c_i$ ) the arrow in  $T_{n,n,n}$  with  $ta_i = x_i, ha_i = x_{i+1}$  (resp.  $tb_i = y_i, hb_i = y_{i+1}, tc_i = z_i, hc_i = z_{i+1}$ ) for  $1 \leq i \leq n - 1$ . The space  $\text{Rep}(T_{n,n,n}, \beta)$  can be identified with

$$\bigoplus_{1 \leq i \leq n-1} (\text{Hom}(V(x_i), V(x_{i+1})) \oplus \text{Hom}(V(y_i), V(y_{i+1})) \oplus \text{Hom}(V(z_i), V(z_{i+1})))$$

where we write  $x_n = y_n = z_n = u$ .

The Cauchy formula [4, §A.1] gives the decomposition of  $K[\text{Rep}(T_{n,n,n}, \beta)]$  as a direct sum over the  $3(n - 1)$ -tuples of partitions

$$((\alpha^i)_{1 \leq i \leq n-1}, (\beta^i)_{1 \leq i \leq n-1}, (\gamma^i)_{1 \leq i \leq n-1})$$

of the summands

$$\bigotimes_{1 \leq i \leq n-1} (S_{\alpha^i} V(x_i) \otimes S_{\alpha^i} V(x_{i+1})^* \otimes S_{\beta^i} V(y_i) \otimes S_{\beta^i} V(y_{i+1})^* \otimes S_{\gamma^i} V(z_i) \otimes S_{\gamma^i} V(z_{i+1})^*).$$

Let us denote  $H = \prod_{1 \leq i \leq n-1} (\text{SL}(V(x_i)) \times \text{SL}(V(y_i)) \times \text{SL}(V(z_i)))$ . Then it follows from the Littlewood-Richardson Rule [4, §A.1] that the summand corresponding to the  $3(n - 1)$ -tuple

$$((\alpha^i)_{1 \leq i \leq n-1}, (\beta^i)_{1 \leq i \leq n-1}, (\gamma^i)_{1 \leq i \leq n-1})$$

contains an  $H$ -invariant if and only if we have for each  $i$ ,  $1 \leq i \leq n - 1$ ,

$$\begin{aligned} (\alpha^i)' &= ((i)^{\sigma(x_i)}, (i - 1)^{\sigma(x_{i-1})}, \dots, 1^{\sigma(x_1)}), \\ (\beta^i)' &= ((i)^{\sigma(y_i)}, (i - 1)^{\sigma(y_{i-1})}, \dots, 1^{\sigma(y_1)}), \\ (\gamma^i)' &= ((i)^{\sigma(z_i)}, (i - 1)^{\sigma(z_{i-1})}, \dots, 1^{\sigma(z_1)}) \end{aligned}$$

for some non-negative numbers  $\sigma(x_i), \sigma(y_i), \sigma(z_i)$ . Moreover, if these conditions are satisfied, then the space of  $H$ -invariants is isomorphic to

$$S_{\alpha^{n-1}} V(u)^* \otimes S_{\beta^{n-1}} V(u)^* \otimes S_{\gamma^{n-1}} V(u)^*.$$

Therefore the space of  $\text{SL}(T_{n,n,n}, \beta)$ -semi-invariants can be identified with the space of  $\text{SL}(V(u))$ -invariants in the above triple tensor product.  $\square$

**Corollary 2.** *The set of triples of partitions  $(\lambda, \mu, \nu)$  such that the space of  $\text{SL}(U)$ -invariants in  $S_\lambda(U) \otimes S_\mu(U) \otimes S_\nu(U)$  is non-zero, in the space of triples of weights is given by a finite set of linear homogeneous inequalities in the parts of  $\lambda, \mu, \nu$  and the condition that  $|\lambda| + |\mu| + |\nu|$  is divisible by  $n := \dim U$ .*

*Proof.* Let  $\sigma \in \Gamma$  be given by (3) and let  $\sigma(\beta) = 0$ . All components of  $\sigma$  are integers only if  $|\lambda| + |\mu| + |\nu|$  is divisible by  $n$ , because

$$0 = \sigma(\beta) = n\sigma(u) + \sum_{i=1}^{n-1} i(\sigma(x_i) + \sigma(y_i) + \sigma(z_i)) = n\sigma(u) + |\lambda| + |\mu| + |\nu|.$$

By Theorem 3 and Proposition 1, those  $(\lambda, \mu, \nu)$  for which  $\text{SI}(T_{n,n,n}, \beta)_\sigma \neq 0$  are given by  $\sigma(\beta) = 0$  and a finite set of homogeneous linear inequalities in  $\sigma(x_i), \sigma(y_i), \sigma(z_i)$ ,  $1 \leq i \leq n - 1$ . These inequalities can be written as inequalities in the parts of  $\lambda, \mu$  and  $\nu$ .  $\square$

#### 4. THE PROOF OF THEOREM 1

We define  $[x, y]$  to be the vector space with the basis formed by paths from  $x$  to  $y$ . We assumed that  $Q$  has no oriented cycles, so the spaces  $[x, y]$  are finite dimensional.

The indecomposable projective representations are in a bijection with  $Q_0$ . The indecomposable projective corresponding to  $x$  is defined by

$$P_x(y) = [x, y], \quad P_x(a) = a \circ \cdot : [x, ta] \rightarrow [x, ha],$$

where  $P_x(a)$  is given by the composition  $p \mapsto a \circ p$ . We have  $\text{Hom}_Q(P_x, V) = V(x)$ . In particular  $\text{Hom}_Q(P_x, P_y) = [y, x]$ .

We choose a numbering  $Q_0 = \{x_1, \dots, x_n\}$  of vertices of  $Q$  such that for every  $\alpha \in Q_1$  with  $t\alpha = x_i, h\alpha = x_j$ , we have  $i < j$ . Let  $b_{i,j}$  be the number of arrows  $\alpha \in Q_1$  with  $t\alpha = x_i, h\alpha = x_j$ . Let  $p_{i,j} = \dim[x_i, x_j]$  be the number of paths  $p$  in  $Q$  such that  $tp = x_i, hp = x_j$ .

The relations between the  $\alpha(x_j)$  and  $\sigma(x_i)$  are as follows:

$$(4) \quad \sigma(x_j) = \alpha(x_j) - \sum_{i < j} b_{i,j} \alpha(x_i),$$

$$(5) \quad \alpha(x_j) = \sigma(x_j) + \sum_{i < j} p_{i,j} \sigma(x_i).$$

We define the  $m$ -arrow quiver  $\Theta_m$  as a quiver with two vertices  $x_+$  and  $x_-$ , and  $m$  arrows  $a_1, \dots, a_m$  with  $ta_i = x_-, ha_i = x_+$  for  $i = 1, \dots, m$ . We define the weight  $\tau$  given by  $\tau(x_+) = 1, \tau(x_-) = -1$ . The dimension vector  $\theta(n)$  is defined by  $\theta(n)(x_+) = \theta(n)(x_-) = n$ .

The idea of the proof of Theorem 1 is to reduce the calculation to the weight space  $\text{SI}(\Theta_m, \theta(n))_\tau$ . The method comes from Classical Invariant Theory with a slight adjustment to accomodate the definition of semi-invariants  $c^V$ .

*Proof of Theorem 1.* Let us fix  $Q, \beta$  and a weight  $\sigma$ . We proceed in three steps. In the first step, we reduce the theorem to the case that  $Q$  is a quiver with exactly one source  $x_-$  and one sink  $x_+$ , and  $\sigma(x_-) = 1, \sigma(x_+) = -1$  and  $\sigma$  is zero on all other vertices. In the second step we reduce to the case that there are no vertices  $x$  with  $\sigma(x) = 0$ . The only case left is the quiver  $\Theta_m$  with weight  $\tau$ . In Step 3 we will prove the theorem in this case.


**Step 1.** Construct a quiver  $Q(\sigma)$  as follows:

$$Q(\sigma)_0 = Q_0 \cup x_- \cup x_+,$$


$$Q(\sigma)_1 = Q_1 \cup Q_- \cup Q_+$$

where  $Q_-$  consists of the set of arrows from  $x_-$  to  $x_i$ , with  $\sigma(x_i)$  arrows going to the vertex  $x_i$  for which  $\sigma(x_i) > 0$  and no arrows going to other vertices. The set  $Q_+$  consists of the set of arrows from  $x_i$  to  $x_+$ , with  $-\sigma(x_i)$  arrows going from the vertex  $x_i$  for which  $\sigma(x_i) < 0$  and no arrows going from other vertices to  $x_+$ .

**Example 2.** Let  $Q$  be the quiver



Let  $\sigma = (1, 1, -2)$ . Then the quiver  $Q(\sigma)$  is



We will write  $\overline{Q} = Q(\sigma)$ . Define the weight  $\overline{\sigma}$  of  $\overline{Q}$  by  $\overline{\sigma}(x_-) = 1, \overline{\sigma}(x_i) = 0, \overline{\sigma}(x_+) = -1$ . The dimension vector  $\overline{\beta} = \beta(\sigma)$  is defined by  $\overline{\beta}(x_i) = \beta(x_i), \overline{\beta}(x_-) = \sum_{\{i|\sigma(x_i)>0\}} \sigma(x_i)\beta(x_i), \overline{\beta}(x_+) = \sum_{\{i|\sigma(x_i)<0\}} -\sigma(x_i)\beta(x_i)$ . Suppose that  $W \in \text{Rep}(\overline{Q}, \overline{\beta})$ . The matrices of all maps  $W(a)$  with  $a \in Q_-$  form a square matrix. Let  $D^-(W)$  be the determinant of this block matrix. Let  $D^+(W)$  be the determinant of all  $W(a)$  with  $a \in Q_+$ . Then the correspondence  $c \rightarrow D^-cD^+$  gives the isomorphism of weight spaces  $\text{SI}(Q, \beta)_\sigma \rightarrow \text{SI}(\overline{Q}, \overline{\beta})_{\overline{\sigma}}$ .

Let  $\overline{\alpha}$  be the dimension vector of  $\overline{Q}$  such that  $\overline{\sigma} = \langle \overline{\alpha}, \cdot \rangle$ . Let  $\overline{V}$  be a representation of  $\overline{Q}$  with dimension vector  $\overline{\alpha}$  and let  $c^{\overline{V}}$  be the corresponding non-zero semi-invariant on  $\text{SI}(\overline{Q}, \overline{\beta})$ .

**Proposition 2.** *The factor  $c$  in the decomposition  $c^{\overline{V}} = D^-cD^+$  is of the form  $c^V$  for some  $V \in \text{Rep}(Q, \alpha)$ .*

*Proof.* Notice that the weight of  $D^-$  is equal to  $\langle \gamma_-, \cdot \rangle$  where

$$\gamma_-(x_-) = 1, \quad \gamma_-(x_j) = \gamma_-(x_+) = 0.$$

Similarly, by (5), the weight of  $D^+$  equals  $\langle \gamma_+, \cdot \rangle$  where

$$\begin{aligned} \gamma_+(x_-) &= 0, \quad \gamma_+(x_j) = - \sum_{\substack{i \leq j \\ \sigma(x_i) < 0}} p_{i,j} \sigma(x_i), \\ \gamma_+(x_+) &= -1 + \sum_{\substack{j \\ \sigma(x_j) < 0}} \sum_{\substack{i \leq j \\ \sigma(x_i) < 0}} p_{i,j} \sigma(x_i) \sigma(x_j). \end{aligned}$$

It is easy to see that  $\langle \gamma_-, \overline{\beta} \rangle = \langle \gamma_+, \overline{\beta} \rangle = 0$ .

Let  $\overline{V} \in \text{Rep}(\overline{Q}, \overline{\alpha})$ . Then  $\overline{V}$  has an obvious submodule  $\overline{V}_1 = \overline{V}|_{\overline{Q}_0 \setminus \{x_-\}}$ . We have an exact sequence

$$0 \rightarrow \overline{V}_1 \rightarrow \overline{V} \rightarrow \overline{V}_2 \rightarrow 0$$

with the dimension of  $\overline{V}_2$  equal to  $\gamma_-$ .

Let  $M$  be the module defined by the exact sequence

$$0 \rightarrow P_{x_+} \xrightarrow{i} \bigoplus_{b, hb=x_+} P_{tb} \rightarrow M \rightarrow 0,$$

where the morphism  $i$  from  $P_{x_+}$  to a copy  $P_{tb}$  maps the trivial path  $e(x_+)$  to the path  $b$ . The dimension vector of  $M$  is  $\gamma_+$ , and  $c^M$  is the determinant  $D^+$ . Consider the map

$$\sum_b \overline{V}_1(b) : \bigoplus_{b, hb=x_+} \overline{V}_1(tb) \rightarrow \overline{V}_1(x_+).$$

The dimension of the kernel is at least 1. Let  $(s_b)_{b, hb=x_+}$  with  $s_b \in \overline{V}_1(tb)$  be a non-trivial element in the kernel. We can now define a map  $\bigoplus_{b, hb=x_+} P_{tb} \rightarrow \overline{V}_1$  by sending the generator  $e(tb) \in P_{tb}(tb)$  to  $s_b$  for all  $b$ . Because  $(s_b)_{b, hb=x_+}$  lies in the kernel, this actually defines a morphism  $M \rightarrow \overline{V}_1$ . Let  $\overline{V}_3$  be the image of this morphism.

Now  $\overline{V}_3$  is a submodule of  $\overline{V}_1$  and  $c^{\overline{V}_3} \neq 0$ . By Lemma 1 a) we have  $\langle \underline{d}(\overline{V}_3), \overline{\beta} \rangle \geq 0$ . We also have  $c^M = D^+ \neq 0$ . If we apply Lemma 1 a) to the kernel  $N$  of

$M \rightarrow \bar{V}_3$ , then we get  $\langle \underline{d}(N), \bar{\beta} \rangle = \langle \gamma_+, -\underline{d}(\bar{V}_3) \rangle = -\langle \underline{d}(\bar{V}_3), \bar{\beta} \rangle \geq 0$ . We conclude that  $\langle \underline{d}(\bar{V}_3), \bar{\beta} \rangle = 0$ . By Lemma 1 b)  $c^{\bar{V}_3}$  divides the semi-invariant  $c^M = D^+$ . Because  $D^+$  is an irreducible semi-invariant we must have  $c^{\bar{V}_3} = D^+$ ,  $\gamma_+ = \dim \bar{V}_3$  and  $\bar{V}_3$  is isomorphic to  $M$ .

We have an exact sequence

$$0 \rightarrow \bar{V}_3 \rightarrow \bar{V}_1 \rightarrow \bar{V}_4 \rightarrow 0.$$

Now it is clear by the multiplicative property that  $c^{\bar{V}} = c^{\bar{V}_2} c^{\bar{V}_4} c^{\bar{V}_3}$  with the first factor being proportional to  $D^-$  and the last one to  $D^+$ . Let us also define a submodule  $\bar{V}_5 = \bar{V}_4|_{\{x_+\}}$ , so we have an exact sequence

$$0 \rightarrow \bar{V}_5 \rightarrow \bar{V}_4 \rightarrow \bar{V}_6 \rightarrow 0.$$

Note that  $\bar{V}_6$  has support within  $Q$ . The restriction of  $\bar{V}_6$  to  $Q$  will be denoted by  $V$ . We will prove that the restriction of  $c^{\bar{V}}$  to  $\text{Rep}(Q, \beta)$  is  $c^V$ .

Extend  $W \in \text{Rep}(Q, \beta)$  to the module  $\bar{W}$  of dimension  $\bar{\beta}$  by putting  $\bar{W}(x_-) = \bigoplus_{a, ta=x_-} W(ha)$ ,  $\bar{W}(x_+) = \bigoplus_{b, hb=x_+} W(tb)$ , with the maps  $\bar{W}(a)$  and  $\bar{W}(b)$  being the components of the identity map. Define the canonical submodule  $\bar{W}_1 = \bar{W}|_{\{x_+\}}$ . We have an exact sequence

$$0 \rightarrow \bar{W}_1 \rightarrow \bar{W} \rightarrow \bar{W}_2 \rightarrow 0.$$

Define the submodule  $\bar{W}_3 = \bar{W}_2|_{\hat{Q} \setminus \{x_-\}}$  of  $\bar{W}_2$ . Now we have an exact sequence

$$0 \rightarrow \bar{W}_3 \rightarrow \bar{W}_2 \rightarrow \bar{W}_4 \rightarrow 0.$$

The representation  $\bar{W}_3$  has support within  $Q$  and its restriction to  $Q$  is just  $W$ .

We now have

$$c^{\bar{V}}(\bar{W}) = c^{\bar{V}_4}(\bar{W}) = c^{\bar{V}_4}(\bar{W}_1) c^{\bar{V}_4}(\bar{W}_3) c^{\bar{V}_4}(\bar{W}_4) = c^{\bar{V}_4}(\bar{W}_3)$$

because  $c^{\bar{V}_4}(\bar{W}_1)$  and  $c^{\bar{V}_4}(\bar{W}_4)$  are constant. Moreover,

$$c^{\bar{V}_4}(\bar{W}_3) = c^{\bar{V}_5}(\bar{W}_3) c^{\bar{V}_6}(\bar{W}_3) = c^{\bar{V}_6}(\bar{W}_3) = c^V(W)$$

because  $c^{\bar{V}_5}(\bar{W}_4)$  is constant. This concludes the proof of the proposition. □

**Step 2.** Let  $Q, \beta, \sigma$  be as above. Let  $x \in Q_0$  be a vertex such that  $\sigma(x) = 0$ . Let  $a_1, \dots, a_s$  be the arrows in  $Q_1$  with  $ha_k = x$  ( $k = 1, \dots, s$ ) and let  $b_1, \dots, b_t$  be the arrows in  $Q_1$  with  $tb_l = x$  ( $l = 1, \dots, t$ ). Let  $\bar{Q}$  be the quiver such that  $\bar{Q}_0 = Q_0 \setminus \{x\}$  and  $\bar{Q}_1 = (Q_1 \setminus \{a_1, \dots, a_s, b_1, \dots, b_t\}) \cup \{ba_{k,l}\}_{1 \leq k \leq s, 1 \leq l \leq t}$ , where  $t(ba_{k,l}) = ta_k, h(ba_{k,l}) = hb_l$ . Let  $\bar{\beta}, \bar{\sigma}$  be the restrictions of  $\beta, \sigma$  to  $Q_0 \setminus \{x\}$ .

The Fundamental Theorem of Invariant Theory (see [2] for a characteristic free version) says that every semi-invariant from  $\text{SI}(Q, \beta)_\sigma$  can be obtained from the semi-invariants from  $\text{SI}(\bar{Q}, \bar{\beta})_{\bar{\sigma}}$  by substituting the actual compositions  $b_l a_k$  for the arrows of type  $ba_{k,l}$ . Assuming Theorem 1 for  $\text{SI}(\bar{Q}, \bar{\beta})_{\bar{\sigma}}$  to be true, we need to show that every semi-invariant  $c^{\bar{V}}$  from  $\text{SI}(\bar{Q}, \bar{\beta})_{\bar{\sigma}}$  pulls back to a semi-invariant of type  $c^V$ . For a given representation  $\bar{V}$  of  $\bar{Q}$  of dimension  $\bar{\alpha}$  we define the representation  $V = \text{ind } \bar{V}$  as follows. We notice that the condition  $\sigma(x) = 0$  means that we expect  $\dim V(x) = \sum_{k=1}^s \dim V(ta_k)$ .

This means we put

$$V(y) = \begin{cases} \overline{V}(y) & \text{if } y \neq x, \\ \bigoplus_{k=1}^s \overline{V}(ta_k) & \text{if } y = x. \end{cases}$$

We define the linear maps  $V(a)$  as follows:

$$V(a) = \begin{cases} \overline{V}(a) & \text{if } a \neq a_k, b_l, \\ i(a_k) & \text{if } a = a_k, \\ \sum_{k=1}^s \overline{V}(ba_{k,l}) & \text{if } b = b_l, \end{cases}$$

where  $i(a_k) : V(ta_k) \rightarrow \bigoplus_{k=1}^s V(ta_k)$  is the injection on the  $k$ -th summand.

Then it is easy to check directly from the definition of semi-invariants  $c^V$  that if the representation  $\overline{W} = \text{res } W$  of dimension  $\overline{\beta}$  is a restriction of a representation  $W$  of  $Q$  of dimension  $\beta$ , then  $c^{\overline{V}}(\overline{W}) = c^V(W)$ .

Notice that the functor  $\text{ind } \overline{V}$  is the left adjoint of the obvious restriction functor  $\text{res} : \text{Rep}(Q) \rightarrow \text{Rep}(\overline{Q})$ , i.e., we have the natural isomorphisms

$$\text{Hom}_Q(\text{ind } \overline{V}, W) = \text{Hom}_{\overline{Q}}(\overline{V}, \text{res } W)$$

which explains why  $c^{\overline{V}}(\overline{W})$  and  $c^V(W)$  vanish simultaneously.

**Step 3.** It remains to deal directly with the weight space  $\text{SI}(\Theta_m, \theta(n))_\tau$ . Writing the representation  $W$  of dimension  $\theta(n)$  as an  $m$ -tuple of linear maps,

$$W(a_1), \dots, W(a_m) : W_- \rightarrow W_+,$$

we can introduce the additional action of the group  $\text{GL}(m)$  acting on this space by taking linear combinations of the linear maps  $W(a_1), \dots, W(a_m)$ . Using the Cauchy formula (in its characteristic free version, say from [1]) we see that the space  $\text{SI}(\Theta_m, \theta(n))_\tau$  of semi-invariants can be identified with  $\bigwedge^n W_- \otimes \bigwedge^n W_+^* \otimes D_n(K^m)$ . Here  $D_n$  denotes the  $n$ -th divided power. Since the divided power  $D_n(K^m)$  is generated as a  $\text{GL}(m)$ -module by its highest weight vector (which corresponds to the semi-invariant  $\det W(a_1)$ ) and the set of semi-invariants of the form  $c^V$  is preserved by the action of  $\text{GL}(m)$ , it is enough to express  $\det W(a_1)$  as the semi-invariant of the form  $c^V$ . Notice that  $\tau = \langle \alpha, \cdot \rangle$  for the dimension vector  $\alpha = (1, m - 1)$ . Taking the module  $V$  to be the  $m$ -tuple of linear maps  $V(a_1), \dots, V(a_m) : K \rightarrow K^{m-1}$  where  $V(a_1) = 0$  and  $V(a_i)$  is the embedding sending 1 to the  $i - 1$ 'st basis vector, for  $i = 2, \dots, m$ , we check directly that  $c^V = \det W(a_1)$ . This concludes the proof of Theorem 1. □

We now will give another description for semi-invariants  $\text{SI}(Q, \beta)_\sigma$ . Let  $\overline{Q} = Q(\sigma), \overline{\beta}$  and  $\overline{\sigma}$  be as in Step 1 of the proof of Theorem 1. We know that  $\text{SI}(Q, \beta)_\sigma \cong \text{SI}(\overline{Q}, \overline{\beta})_{\overline{\sigma}}$ . Let  $\overline{\alpha}$  be a dimension vector of  $\overline{Q}$  such that  $\langle \overline{\alpha}, \cdot \rangle = \overline{\sigma}$ . Now  $\text{SI}(\overline{Q}, \overline{\beta})_{\overline{\sigma}}$  is generated by semi-invariants  $c^{\overline{V}}$  with  $\underline{d}(\overline{V}) = \overline{\alpha}$ . In fact we only need to take those  $c^{\overline{V}}$  where  $\overline{V}$  lies in a Zariski dense set of  $\text{Rep}(\overline{Q}, \overline{\alpha})$ . A general representation  $\overline{V}$  of dimension  $\overline{\alpha}$  has the following projective resolution:

$$0 \rightarrow P_{x_+} \xrightarrow{d_V} P_{x_-} \rightarrow \overline{V} \rightarrow 0$$

with  $d_V \in \text{Hom}_Q(P_{x_+}, P_{x_-}) = [x_-, x_+]$ . So  $d_V$  can be seen as some linear combination  $\sum_{i=1}^r \lambda_i p_i$  where  $p_1, \dots, p_r$  are all paths from  $x_+$  to  $x_-$ . For any  $\overline{W} \in \text{Rep}(\overline{Q}, \overline{\beta})$  we have the following exact sequence:

$$0 \rightarrow \text{Hom}_{\overline{Q}}(\overline{V}, \overline{W}) \rightarrow \text{Hom}_{\overline{Q}}(P_{x_+}, \overline{W}) \xrightarrow{\tilde{d}_{\overline{V}}} \text{Hom}_{\overline{Q}}(P_{x_-}, \overline{W}) \rightarrow \text{Ext}_{\overline{Q}}(\overline{V}, \overline{W}) \rightarrow 0.$$

It is easy to see that  $\det(\tilde{d}_{\overline{V}}) = c^{\overline{V}}(\overline{W}) = c^V(W)$ .

We have that

$$\begin{aligned}\mathrm{Hom}_{\overline{Q}}(P_{x_+}, \overline{W}) &\cong \overline{W}_{x_+} = \bigoplus_{\sigma(x_i) > 0} W(x_i)^{\sigma(x_i)}, \\ \mathrm{Hom}_{\overline{Q}}(P_{x_-}, \overline{W}) &\cong \overline{W}_{x_-} = \bigoplus_{\sigma(x_i) < 0} W(x_i)^{\sigma(x_i)}, \\ \tilde{d}_{\overline{V}} &= \sum_i \lambda_i \overline{V}(p_i).\end{aligned}$$

Let  $F$  be a function from the set of paths from  $x_+$  to  $x_-$  to the set of non-negative integers. For each such  $F$  we can define the semi-invariant  $I_F$  as the coefficient of  $\lambda_1^{F(p_1)} \lambda_2^{F(p_2)} \dots \lambda_r^{F(p_r)}$  in  $\det(\tilde{d}_{\overline{V}})$ .

**Corollary 3.** *The space of semi-invariants  $\mathrm{SI}(Q, \beta)_\sigma$  is spanned by semi-invariants of the form  $I_F$ .*

A necessary condition for  $I_F$  to be non-zero is

$$\sum_i F(p_i) = \sum_{\sigma(x_i) > 0} \sigma(x_i) \beta(x_i) = \sum_{\sigma(x_i) < 0} -\sigma(x_i) \beta(x_i).$$

#### ACKNOWLEDGEMENT

The authors would like to thank Andrei Zelevinsky for helpful suggestions.

#### REFERENCES

- [1] K. Akin, D. A. Buchsbaum, J. Weyman, *Schur functors and Schur complexes*, Adv. Math. **44** (1982), 207–278. MR **84c**:20021
- [2] C. DeConcini, C. Procesi, *Characteristic free approach to invariant theory*, Adv. Math. **21** (1976), 330–354. MR **54**:10305
- [3] W. Fulton, *Eigenvalues of sums of Hermitian matrices* (after A. Klyachko), Séminaire Bourbaki (1998). MR **99m**:00026
- [4] W. Fulton, J. Harris, *Representation Theory*, Springer-Verlag, New York, 1991. MR **93a**:20069
- [5] V. Kac, *Infinite root systems, representations of graphs and invariant theory II*, J. Algebra **78** (1982), 141–162. MR **85b**:17003
- [6] A. D. King, *Moduli of representation of finite dimensional algebras*, Quart. J. Math. Oxford (2) **45** (1994), 515–530. MR **96a**:16009
- [7] A. Klyachko, *Stable vector bundles and Hermitian operators*, IGM, University of Marne-la-Vallée, preprint (1994).
- [8] A. Knutson, T. Tao, *The honeycomb model of  $\mathrm{GL}_n(\mathbb{C})$  tensor products, I: Proof of the saturation conjecture*, J. Amer. Math. Soc. **12** (1999), 1055–1090. MR **2000c**:20066
- [9] C.M. Ringel, *Representations of  $K$ -species and bimodules*, J. Algebra **41** (1976) 269–302. MR **54**:10340
- [10] A. Schofield, *Semi-invariants of quivers*, J. London Math. Soc. **43** (1991), 383–395. MR **92g**:16019
- [11] A. Schofield, *General representations of quivers*, Proc. London Math. Soc. (3) **65** (1992) 46–64. MR **93d**:16014
- [12] A. Schofield, M. van den Bergh, *Semi-invariants of quivers for arbitrary dimension vectors*, preprint, math.RA/9907174.

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MASSACHUSETTS 02151

*E-mail address:* hderksen@math.mit.edu

DEPARTMENT OF MATHEMATICS, NORTHEASTERN UNIVERSITY, BOSTON, MASSACHUSETTS 02115

*E-mail address:* weyman@neu.edu